

A liquid bait for the reduction of reproductive capacity in rats.

For indoor use in rafters, ceilings, attics, and other elevated areas of agricultural farm buildings, transit systems, public and private buildings, commercial residential buildings, retail buildings, and/or other man-made structures.

Not Intended for Consumer Use

## **Active Ingredients:**

4-Vinylcyclohexene diepoxide	0.09604%
Triptolide	0.00118%
Other Ingredients:	99.90278%
•	Total: 100 0000%

# Keep Out of Reach of Children **CAUTION**

Made in USA EPA Reg. No. 91601-1 EPA Est. No. 91601-AZ-1 Net Contents: 1-20 fl. oz. (30-592 ml)

## FIRST AID

#### If on skin or clothing:

- · Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15–20 minutes.
- · Call a poison control center or doctor for treatment advice.

#### Note to Physician or Veterinarian:

This product has been formulated for fertility control in rats. It may possibly induce transient infertility in small male and female mammals capable of reproduction.

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

## PRECAUTIONARY STATEMENTS Hazards to Humans and Domestic Animals

**CAUTION.** Avoid contact with skin or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

## **Personal Protective Equipment:**

Wear protective nitrile gloves ≥ 14 mils or butyl rubber ≥ 14 mils, long sleeved shirts, and long pants when handling this product.

**Treatment for Pet Ingestion:** If animal consumes bait, and adverse health effects are observed, call a veterinarian.

#### DIRECTIONS FOR USE

<u>READ THIS LABEL</u>: Read label attached to product. Follow all use directions and use precautions. Use only for sites, pests, and application methods described on this label. It is a violation of Federal law to use this product in a manner inconsistent with its labeling. <u>FOR INDOOR NON-FOOD USE</u>: in rafters, ceilings, attics, and other elevated areas of agricultural farm buildings, transit systems, public and private buildings, commercial residential buildings, retail buildings, and/or other man-made structures. DO NOT apply bottles outdoors.

DO NOT apply bottles in areas accessible to children, pets or non-target species.

## **USE RESTRICTIONS:**

- 1. Bottles must not be placed in areas where the contamination of food or surfaces that come in direct contact with food is possible.
- 2. Place bottles indoors only, DO NOT apply outdoors.
- 3. Do not dilute or add anything to the product.
- 4. Store product in a location where children, pets, or other non-target animals have no access.
- 5. Dispose of product container, unused, spoiled, and unconsumed bait as specified on this label.
- 6. Bottle mounting bracket is mandatory for use. Mounting bracket may either be secured to a sturdy vertical or horizontal surface. <u>SELECTION OF TREATMENT AREAS</u>: Determine areas where rats will most likely find and consume bait. Generally these are along ceiling rafters, ledges, or beams, attics, along walls, or in locations where rats or their signs have been observed.

APPLICATION DIRECTIONS: Maintain an uninterrupted supply of bait. Check and replace bait as necessary, or at least every month. Installing Bottles into Mounting Bracket:

# To use bottle and mounting bracket:

- 1. Secure mounting bracket to a solid surface. When placed on a vertical surface, secure the bracket so that it does not move side to side. When placed on horizontal surface, secure the bracket so that it cannot be knocked over.
- 2. Adjust the height of the bracket to ensure that once the bottle and nozzle are attached, rats can consume from the drinking nozzle.
- 3. Remove the cap and seal from the bottle.
- 4. Attach the nozzle to the bottle.
- 5. Invert the bottle and place in the mounting bracket.
- 6. Once inverted and in place, remove the stopper covering the end of the nozzle.
- 7. It is recommended to place a secondary container under mounting bracket when hung from a vertical surface.

## Installing replacement bottle:

- 1. Remove the bottle from the mounting bracket by pulling straight up. **Note:** If you are unable to grip the bottle securely to remove, you can push the bottle up from underneath.
- 2. Invert the bottle.
- 3. Remove the nozzle and recap the empty bottle with the cap from the replacement bottle and dispose of (see storage and disposal). If necessary, replace the nozzle.
- 4. Remove the cap and seal from the replacement bottle.
- 5. Attach the drinking nozzle to the replacement bottle.
- 6. Invert the replacement bottle and place the bottle in the mounting bracket.
- 7. Once inverted and in place, remove the plug covering the end of the drinking nozzle.

#### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**Storage:** Store only in original, tightly secured, bottles in a room temperature environment, inaccessible to unauthorized users, children, pets, livestock, and other non-target animals. **Disposal:** Do not re-use empty bottles. **If empty:** Place container in trash or offer for recycling. **If partly filled:** Cap the bottle and place in the trash. **Container Handling:** Bottles are not to be reused or refilled. Dispose of bottles by recapping and placing in the trash or offer for recycling at approved waste disposal facility.

#### DISCLAIMER

To the extent consistent with applicable law, seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

Patent 9,956,235 Manufactured by: SenesTech, Inc., 777 W Pinnacle Peak Rd, Suite B104, Phoenix, AZ 85027

ContraPest® is a registered trademark of SenesTech, Inc.